

ONE LEVEL DENSITY OF LOW-LYING ZEROS OF FAMILIES OF L -FUNCTIONS

PENG GAO AND LIANGYI ZHAO

ABSTRACT. In this paper, we prove some one level density results for the low-lying zeros of families of L -functions. More specifically, the families under consideration are that of L -functions of holomorphic Hecke eigenforms of level 1 and weight k twisted with quadratic Dirichlet characters and that of cubic and quartic Dirichlet L -functions.

Mathematics Subject Classification (2010): 11F11, 11L40, 11M06, 11M26, 11M50, 11R16

Keywords: one level density, low-lying zeros, Hecke eigenforms, quadratic Dirichlet characters, cubic Dirichlet characters, quartic Dirichlet characters

CONTENTS

1. Introduction	1
1.5. Notations	5
2. Preliminaries	5
2.1. The Explicit Formula	5
2.5. Poisson Summation	7
2.8. Primitive cubic and quartic Dirichlet characters	8
3. Proof of Theorem 1.1	9
3.1. Estimation of $S_R(X, Y; \hat{\phi}, f, \Phi)$	9
3.3. Estimation of $S_M(X, Y; \hat{\phi}, f, \Phi)$	9
3.4. Conclusion	10
4. Proof of Theorem 1.2	10
5. Proof of Theorem 1.3 and Theorem 1.4	13
6. Notes	14
Acknowledgments	14
References	14

1. INTRODUCTION

The density conjecture of N. Katz and P. Sarnak [21, 22] suggests that the distribution of zeros near the central point of a family of L -functions is the same as that of eigenvalues near 1 of a corresponding classical compact group. This has been confirmed for many families of L -functions, such as different types of Dirichlet L -functions [10, 15, 26, 27, 30], L -functions with characters of the ideal class group of the imaginary quadratic field $\mathbb{Q}(\sqrt{-D})$ [9], automorphic L -functions [8, 14, 19, 28, 29], elliptic curves L -functions [2, 3, 13, 25, 34], symmetric powers of $GL(2)$ L -functions [7, 12] and a family of $GL(4)$ and $GL(6)$ L -functions [7]. Literatures in this direction are too numerous to be completely listed here. In this paper, we prove some one level density results for the low-lying zeros of families of L -functions of holomorphic Hecke eigenforms of level 1 and weight k twisted with quadratic Dirichlet characters and those of families of cubic and quartic Dirichlet L -functions.

Let χ be a primitive Dirichlet character and we denote the non-trivial zeroes of the Dirichlet L -function $L(s, \chi)$ by $\frac{1}{2} + i\gamma_{\chi, j}$. Without assuming the generalized Riemann hypothesis (GRH), we order them as

$$(1.1) \quad \dots \leq \Re\gamma_{\chi, -2} \leq \Re\gamma_{\chi, -1} < 0 \leq \Re\gamma_{\chi, 1} \leq \Re\gamma_{\chi, 2} \leq \dots$$

For any primitive Dirichlet character χ of conductor q of size X , we set

$$\tilde{\gamma}_{\chi, j} = \frac{\gamma_{\chi, j}}{2\pi} \log X$$

and define for an even Schwartz class function ϕ ,

$$(1.2) \quad S(\chi, \phi) = \sum_j \phi(\tilde{\gamma}_{\chi, j}).$$

In [27], A. E. Özlük and C. Snyder studied the family of quadratic Dirichlet L -functions. Consider the family of Dirichlet L -functions of the form $L(\chi_{8d}, s)$ for d odd and square-free with $X \leq d \leq 2X$, where $\chi_{8d} = (\frac{8d}{\cdot})$ is the Kronecker symbol. Let $D(X)$ denote the set of such d 's. It's easy to see, as noted in [10], that

$$\#D(X) \sim \frac{4X}{\pi^2}.$$

Assuming GRH for this family, it follows from the result of Özlük and Snyder [27] that for $W_{USp}(x) = 1 - \frac{\sin(2\pi x)}{2\pi x}$, we have

$$(1.3) \quad \frac{1}{\#D(X)} \sum_{d \in D(X)} S(\chi_{8d}, \phi) \sim \int_{\mathbb{R}} \phi(x) W_{USp}(x) dx,$$

as $X \rightarrow \infty$ provided that the support of $\hat{\phi}$, the Fourier transform of ϕ , is contained in the interval $(-2, 2)$. The expression on the left-hand side of (1.3) is known as the one-level density of the low-lying zeros for this family of L -functions under consideration.

The kernel of the integral W_{USp} in (1.3) is the same function which occurs on the random matrix theory side, when studying the eigenvalues of unitary symplectic matrices. This shows that the family of quadratic Dirichlet L -functions is a symplectic family. In [30], M. O. Rubinstein studied all the n -level densities of the low-lying zeros of the families of quadratic twists of L -functions attached to a self-contragredient automorphic cuspidal representation, as well as the family of quadratic Dirichlet L -functions. He showed that they converge to the symplectic densities for test functions $\phi(x_1, \dots, x_n)$ whose Fourier transforms $\hat{\phi}(u_1, \dots, u_n)$ have their supports contained in the set

$$\left\{ (u_1, \dots, u_n) \in \mathbb{R}^n : \sum_{i=1}^n |u_i| < 1 \right\}.$$

The result of Rubinstein does not assume GRH. In [10], assuming the truth of GRH, the first-named author extended Rubinstein's result [30] and showed that it holds for $\hat{\phi}(u_1, \dots, u_n)$ with support in the set

$$\left\{ (u_1, \dots, u_n) \in \mathbb{R}^n : \sum_{i=1}^n |u_i| < 2 \right\}.$$

In this paper, we consider a few other families of L -functions. First, let f be a fixed holomorphic Hecke eigenform of level 1 and weight k . For $\Im(z) > 0$ we have a Fourier expansion of f :

$$f(z) = \sum_{n=1}^{\infty} a_f(n) n^{\frac{k-1}{2}} e(nz),$$

where the coefficients $a_f(n)$ are real and normalized with $a_f(1) = 1$ and satisfy the Ramanujan-Petersson bound

$$|a(n)| \leq d(n),$$

with $d(n)$ being the divisor function. Let χ be a primitive Dirichlet character of conductor q . The L -function of the twist of f by χ is given by

$$(1.4) \quad L(f \times \chi, s) = \sum_{n=1}^{\infty} \frac{a_f(n)\chi(n)}{n^s} = \prod_p \left(1 - \frac{a_f(p)\chi(p)}{p^s} + \frac{\chi(p^2)}{p^{2s}} \right)^{-1}$$

for $\Re(s) > 1$. This L -function continues to an entire function and satisfies the functional equation

$$\left(\frac{q}{2\pi}\right)^s \Gamma\left(\frac{k}{2} + s\right) L\left(f, \frac{1}{2} + s\right) = \iota_\chi \left(\frac{q}{2\pi}\right)^{-s} \Gamma\left(\frac{k}{2} - s\right) L\left(f \times \bar{\chi}, \frac{1}{2} - s\right).$$

where $\iota_\chi = i^k \tau(\chi)^2/q$ and $\tau(\chi)$ is the Gauss sum associated to χ (thus $|\iota_\chi| = 1$). cf. [18, Prop. 14.20]. See [17, 18] for detailed discussions of Hecke eigenforms.

For a fixed f , we consider the family of quadratic twists of L -functions $L(f \times \chi_{8d}, s)$ for d odd and square-free with $X \leq d \leq 2X$. For this family, Rubinstein [30] has shown that the n -level densities of the low-lying zeros converge to the orthogonal densities for test functions $\phi(x_1, \dots, x_n)$ whose Fourier transforms $\hat{\phi}(u_1, \dots, u_n)$ are supported in the set

$$\left\{ (u_1, \dots, u_n) \in \mathbb{R}^n : \sum_{i=1}^n |u_i| < \frac{1}{2} \right\}.$$

More precisely, we denote the zeroes of $L(f \times \chi_{8d}, s)$ by $\frac{1}{2} + i\gamma_{f,8d,j}$, and order them in a manner similar to (1.1). Let $\tilde{\gamma}_{f,8d,j} = \gamma_{f,8d,j} 2 \log X / (2\pi)$ and define for an even Schwartz function ϕ ,

$$D(d, f, \phi) = \sum_j \phi(\tilde{\gamma}_{f,8d,j}).$$

We set

$$W_{SO+}(x) = 1 + \frac{\sin(2\pi x)}{2\pi x}, \quad W_{SO-}(x) = \delta(x) + 1 - \frac{\sin(2\pi x)}{2\pi x} \quad \text{and} \quad W_O(x) = \frac{1}{2} (W_{SO+}(x) + W_{SO-}(x)),$$

where $\delta_0(x)$ is the Dirac distribution at $x = 0$. Note that the three orthogonal densities are indistinguishable for test functions whose Fourier transforms are supported in $(-1, 1)$. The result of Rubinstein asserts that

$$\frac{1}{\#D(X)} \sum_{d \in D(X)} D(d, f, \phi) \sim \int_{\mathbb{R}} \phi(x) \left(1 + \frac{\sin(2\pi x)}{2\pi x} \right) dx,$$

as $X \rightarrow \infty$ provided that $\hat{\phi}$ is supported on the interval $(-1/2, 1/2)$. This shows that the family $L(f \times \chi_{8d}, s)$ has orthogonal symmetry.

In this paper, we improve the above-mentioned result of Rubinstein by doubling the size of the allowable support of the Fourier transform of the test function in the case of the one level density. For technical reasons, we consider the average over the family by a smooth weight. Let $\Phi_X(t)$ be a non-negative smooth function supported on $(1, 2)$, satisfying $\Phi_X(t) = 1$ for $t \in (1 + 1/U, 2 - 1/U)$ with $U = \log X$ and such that $\Phi_X^{(j)}(t) \ll_j U^j$ for all integers $j \geq 0$. Our result is

Theorem 1.1. *Suppose that GRH is true. Let f be a fixed holomorphic Hecke eigenform of level 1 and weight k . Let $\phi(x)$ be an even Schwartz function whose Fourier transform $\hat{\phi}(u)$ has compact support in $(-1, 1)$. Then we have*

$$(1.5) \quad \lim_{X \rightarrow +\infty} \frac{1}{\#D(X)} \sum_{d \in D(X)} \Phi_X\left(\frac{d}{X}\right) D(d, f, \phi) = \int_{\mathbb{R}} \phi(x) \left(1 + \frac{\sin(2\pi x)}{2\pi x} \right) dx.$$

We shall describe briefly the proof of Theorem 1.1 in this paragraph. Using a modified version of the explicit formula, Lemma 2.2, $D(d, f, \phi)$ in (1.5), a sum over the zeros of the L -function under consideration, is converted into a sum over primes and prime powers. The most important among these will be the sums over primes (the one appearing in (2.3)) and prime squares, after showing that the contributions of the higher prime powers are negligible. The sum involving prime squares can be easily handled. Thus far,

each of the sums can be disposed of for each individual L -functions in question, without appealing to the additional averaging over d in (1.5). But this additional sum over the family is needed in the treatment of the more difficult sum involving primes, (2.3). Following a method of K. Soundararajan [32], after detecting the square-free condition of d using the Möbius function and applying Poisson summation formula to the sum over d , we are led to consider sums involving quadratic Gauss sums, which is further split into pieces according to the size of the relevant parameters. GRH is needed to estimate a component of these sums. Other estimates are quoted from [26] to give the final estimate for the sum in (2.3). Theorem 1.1 follows after combining all estimates.

We note that the one level density of the low-lying zeroes of the family of quadratic twists of Hecke L -functions evaluated in Theorem 1.1 is insufficient to determine which of the orthogonal symmetry types, $SO(\text{even})$, O or $SO(\text{odd})$, is attached of each family, due to the small support-restriction on the Fourier transforms of test functions. However, as shown in [7, 25], the two level density allows one to distinguish between the three orthogonal symmetries for test functions whose Fourier transforms have arbitrarily small support. In fact, it follows from Rubinstein's result on two-level density of this family (Lemma 7 in [30] and also the formula on the top of page 179 in [30]) that the symmetry type attached to the family of quadratic twists of L -functions $L(f \times \chi_{8d}, s)$ for d odd and square-free with $X \leq d \leq 2X$ is $SO(\text{even})$.

In addition, we consider the one level density of the low-lying zeros of the families of cubic and quartic Dirichlet L -functions in this paper.

For a primitive cubic Dirichlet character χ of conductor q coprime to 3, it is shown in [1] that q must be square-free and a product of primes congruent to 1 modulo 3. It follows that χ is a product of primitive cubic characters modulo the prime divisors of q and for each prime divisor p of q , there are exactly two primitive characters with conductor p , each being the square (also the complex conjugate) of the other.

We shall prove the following:

Theorem 1.2. *Let $f(x)$ be an even Schwartz function whose Fourier transform $\hat{f}(u)$ has compact support in $(-3/7, 3/7)$, then*

$$\lim_{X \rightarrow +\infty} \frac{1}{\#C(X)} \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* S(\chi, f) = \int_{\mathbb{R}} f(x) dx.$$

Here the “ $*$ ” on the sum over χ means that the sum is restricted to primitive characters and $C(X)$ denotes the set of primitive cubic characters of conductor q not divisible by 3 and $X \leq q \leq 2X$.

We point out here that Theorem 1.2 is analogous to a result of A. M. Güloğlu [11] and that his result depends on GRH while Theorem 1.2 does not. Güloğlu [11] studied the one level density of the low-lying zeros of a family of Hecke L -functions associated with cubic symbols $\chi_c = (\frac{\cdot}{c})_3$ with c square-free and congruent to 1 modulo 9, regarded as primitive ray class characters of the ray class group $h_{(c)}$. We recall here that for any c , the ray class group $h_{(c)}$ is defined to be $I_{(c)}/P_{(c)}$, where $I_{(c)} = \{\mathcal{A} \in I, (\mathcal{A}, (c)) = 1\}$ and $P_{(c)} = \{(a) \in P, a \equiv 1 \pmod{c}\}$ with I and P denoting the group of fractional ideals in $K = \mathbb{Q}(\omega)$ and the subgroup of principal ideals, respectively. Here $\omega = \exp(2\pi i/3)$. The Hecke L -function associated with χ_c is defined for $\Re(s) > 1$ by

$$L(s, \chi_c) = \sum_{0 \neq \mathcal{A} \subset \mathcal{O}_K} \chi_c(\mathcal{A}) (N(\mathcal{A}))^{-s},$$

where $\mathcal{A}$ runs over all non-zero integral ideals in K and $N(\mathcal{A})$ is the norm of $\mathcal{A}$. As shown by E. Hecke, $L(s, \chi_c)$ admits analytic continuation to an entire function and satisfies a functional equation. We refer the reader to [11, 24] for a more detailed discussion of these Hecke characters and L -functions. We denote non-trivial zeroes of $L(s, \chi_c)$ by $\frac{1}{2} + i\gamma_{\chi_c, j}$ and order them in a fashion similar to (1.1). Let $C_{(9)}(X)$ stand for the set of χ_c 's with c square-free, congruent to 1 modulo 9 and $X \leq N(c) \leq 2X$. We define $S(\chi_c, f)$ similarly as $S(\chi, \phi)$ in (1.2).

For the family of Hecke L -functions considered by Güloğlu [11], we can apply our approach to Theorem 1.2 to obtain the following result without assuming GRH:

Theorem 1.3. *Let $f(x)$ be an even Schwartz function whose Fourier transform $\hat{f}(u)$ has compact support in $(-3/5, 3/5)$, then*

$$\lim_{X \rightarrow +\infty} \frac{1}{\#C_{(9)}(X)} \sum_{\substack{c \equiv 1 \pmod{9} \\ X \leq N(c) \leq 2X}}^* S(\chi_c, f) = \int_{\mathbb{R}} f(x) dx.$$

Here the “*” on the sum over c means that the sum is restricted to square-free elements c of $\mathbb{Z}[\omega]$.

The support obtained under GRH by Güloğlu [11] is $(-31/30, 31/30)$ in place of $(-3/5, 3/5)$ in our unconditional Theorem 1.3. In addition, the result Güloğlu’s result is a smoothed version of one-level density and depends on other results that require this smoothness. We also note that, as

$$\int_{\mathbb{R}} f(x) dx = \int_{\mathbb{R}} f(x) W_U(x) dx, \text{ with } W_U(x) = 1,$$

Theorems 1.2 and 1.3 show that the family of cubic Dirichlet L -functions as well as the family of Hecke L -functions associated with cubic symbols are unitary families, an observation made in [11].

Analogous to Theorem 1.2, we also study the one level density of the low-lying zeros of the family of quartic Dirichlet L -functions. We have the following

Theorem 1.4. *Let $f(x)$ be an even Schwartz function whose Fourier transform $\hat{f}(u)$ has compact support in $(-3/7, 3/7)$, then*

$$\lim_{X \rightarrow +\infty} \frac{1}{\#Q(X)} \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^4 = \chi_0, \chi^2 \neq \chi_0}}^* S(\chi, f) = \int_{\mathbb{R}} f(x) dx.$$

Here the “*” on the sum over χ means that the sum is restricted to primitive characters and $Q(X)$ denotes the set of primitive complex quartic characters with odd conductor q and $X \leq q \leq 2X$.

The proofs of Theorems 1.2, 1.3 and 1.4 are similar. Therefore, we describe briefly here only the proof of Theorem 1.2. Similar to the proof of Theorem 1.1, we first convert the sum over zeros of the L -functions under consideration into sums over primes using the relevant versions of the explicit formula. This leads us to consider sums involving the cubic or quartic symbols over primes and prime squares. These are then rewritten as sums involving Hecke characters with new summation conditions. The summation conditions are handled using Möbius inversion and the final estimates are obtained using a Pólya-Vinogradov type bound for these Hecke characters, Lemma 4.1.

1.5. Notations. The following notations and conventions are used throughout the paper.

$e(z) = \exp(2\pi iz) = e^{2\pi iz}$.

$f = O(g)$ or $f \ll g$ means $|f| \leq cg$ for some unspecified positive constant c .

For $x \in \mathbb{R}$, $\|x\| = \min_{n \in \mathbb{Z}} |x - n|$ denotes the distance between x and the closest integer.

2. PRELIMINARIES

2.1. The Explicit Formula. Our approach in this paper relies on the following explicit formula, which essentially converts the sum over zeros of an L -function to the sum over primes.

Lemma 2.2. *Let $f(x)$ be an even Schwartz function whose Fourier transform $\hat{f}(u)$ is compactly supported. Then for any primitive Dirichlet character χ , we have*

$$(2.1) \quad \begin{aligned} S(\chi, f) &= \int_{-\infty}^{\infty} f(t) dt - \frac{1}{\log X} \sum_p \frac{\log p}{\sqrt{p}} \hat{f}\left(\frac{\log p}{\log X}\right) (\chi(p) + \bar{\chi}(p)) \\ &\quad - \frac{1}{\log X} \sum_p \frac{\log p}{p} \hat{f}\left(\frac{2 \log p}{\log X}\right) (\chi(p^2) + \bar{\chi}(p^2)) + O\left(\frac{1}{\log X}\right). \end{aligned}$$

Proof. We combine [31, (2.16)], the fact that $f(x)$ is rapidly decreasing and the Stirling formula which gives that $\Gamma'/\Gamma(s) = \log s + O(1/|s|)$, uniformly for $|\arg s| \leq \pi - \delta$, $|s| \geq 1$ to replace the Γ'/Γ -terms in [31, (2.16)] by $O(1/\log X)$. Moreover, the terms $n = p^k$ for $k \geq 3$, p prime in the sum on the right-hand side of [31, (2.16)] contribute

$$\ll \sum_{p^k, k \geq 3} \frac{\log p}{p^{k/2}} \ll 1.$$

The lemma follows from these observations. $\square$

Similarly, we have [11, Lemma 4.1]:

Lemma 2.3. *Let $f(x)$ be an even Schwartz function whose Fourier transform $\hat{f}(u)$ has compact support. Then for any square-free $c \equiv 1 \pmod{9}$ of $\mathbb{Z}[\omega]$, we have*

$$\begin{aligned} S(\chi_c, f) &= \int_{-\infty}^{\infty} f(t) dt - \frac{1}{\log X} \sum_{\mathfrak{p}} \frac{\log N(\mathfrak{p})}{\sqrt{N(\mathfrak{p})}} \hat{f}\left(\frac{\log N(\mathfrak{p})}{\log X}\right) (\chi_c(\mathfrak{p}) + \overline{\chi}_c(\mathfrak{p})) \\ &\quad - \frac{1}{\log X} \sum_{\mathfrak{p}} \frac{\log N(\mathfrak{p})}{N(\mathfrak{p})} \hat{f}\left(\frac{2 \log N(\mathfrak{p})}{\log X}\right) (\chi_c(\mathfrak{p}^2) + \overline{\chi}_c(\mathfrak{p}^2)) + O\left(\frac{1}{\log X}\right), \end{aligned}$$

where $\mathfrak{p}$ runs over all non-zero integral ideals in $\mathbb{Q}(\omega)$.

For $\Re(s) > 1$, we can rewrite the Euler product (1.4) of $L(f \times \chi_{8d}, s)$ as

$$L(f \times \chi_{8d}, s) = \prod_p \left(1 - \frac{\alpha_f(p) \chi_{8d}(p)}{p^s}\right)^{-1} \left(1 - \frac{\alpha_f^{-1}(p) \chi_{8d}(p)}{p^s}\right)^{-1},$$

with $\alpha_f(p) + \alpha_f^{-1}(p) = a_f(p)$. Similar to Lemma 2.2, we have the following

Lemma 2.4. *Let $\phi(x)$ be an even Schwartz function whose Fourier transform $\hat{\phi}(u)$ is compactly supported. Then for $d \in D(X)$, we have*

$$(2.2) \quad D(d, f, \phi) = \int_{-\infty}^{\infty} \phi(t) dt + \frac{1}{2} \int_{-\infty}^{\infty} \hat{\phi}(u) du - S(f, d, X; \hat{\phi}) + O\left(\frac{\log \log 3X}{\log X}\right),$$

with the implicit constant depending on ϕ and

$$(2.3) \quad S(f, d, X; \hat{\phi}) = \frac{1}{\log X} \sum_p \frac{a_f(p) \log p}{\sqrt{p}} \left(\frac{8d}{p}\right) \hat{\phi}\left(\frac{\log p}{2 \log X}\right).$$

Proof. We use [31, (2.16)] again here and identify $c_\pi(n)$ in [31, (2.16)] as $\Lambda(n) \chi_{8d}(n) a_f(n)$ with $a_f(p^k) = \alpha_f^k(p) + \alpha_f^{-k}(p)$ in our case. By Deligne's proof [6] of the Weil conjecture, we know that $|\alpha_f(p)| = 1$ so that the terms $n = p^k$ for $k \geq 3$, p prime in the sum on the right-hand side of [31, (2.16)] contribute $O(1)$. Moreover, note that

$$\sum_{p|8d} \frac{\log p}{p} \ll \log \log 3X.$$

Thus the terms $n = p^2$ in the sum on the right-hand side of [31, (2.16)] contribute

$$\frac{1}{\log X} \sum_p \frac{a_f(p^2) \log p}{p} \hat{\phi}\left(\frac{\log p}{\log X}\right) + O\left(\frac{\log \log 3X}{\log X}\right).$$

Recall that

$$(2.4) \quad a_f(p^2) = a_f^2(p) - 2$$

and it follows from Proposition 2.3 in [31] that

$$(2.5) \quad \sum_{p \leq x} \frac{a_f^2(p) \log^2 p}{p} = \frac{\log^2 x}{2} + O(\log x).$$

Note also that we have Mertens' formula [4, p. 57],

$$(2.6) \quad \sum_{p \leq x} \frac{\log p}{p} = \log x + O(1).$$

Combining (2.4), (2.5) and (2.6), we deduce that

$$\sum_{p \leq x} \frac{a_f(p^2) \log^2 p}{p} = -\frac{\log^2 x}{2} + O(\log x).$$

From the above and partial summation, we get that

$$\begin{aligned} \frac{1}{\log X} \sum_p \frac{a_f(p^2) \log p}{p} \hat{\phi} \left(\frac{\log p}{\log X} \right) &= -\frac{1}{\log X} \int_1^\infty \hat{\phi} \left(\frac{\log t}{\log X} \right) \frac{dt}{t} + O \left(\frac{\log \log X}{\log X} \right) \\ &= -\frac{1}{2} \int_{-\infty}^\infty \hat{\phi}(t) dt + O \left(\frac{\log \log X}{\log X} \right). \end{aligned}$$

The assertion of the lemma follows from this easily. $\square$

2.5. Poisson Summation. We now fix f and X and henceforth write $\Phi(t)$ in place of $\Phi_X(t)$. We shall focus on finding the asymptotic expression of (with $n \geq 1$)

$$S(X, Y; \hat{\phi}, f, \Phi) := \sum_{\gcd(d, 2)=1} \mu^2(d) \sum_{p \leq Y} \frac{a_f(p) \log p}{\sqrt{p}} \left(\frac{8d}{p} \right) \hat{\phi} \left(\frac{\log p}{2 \log X} \right) \Phi \left(\frac{d}{X} \right),$$

where $\hat{\phi}(u)$ is smooth and has its support contained in the interval $(-1 + \epsilon, 1 - \epsilon)$ for any $\epsilon > 0$. To emphasize this condition, here and throughout, we shall set $Y = X^{2-2\epsilon}$ and write the condition $p \leq Y$ explicitly.

Let $Z > 1$ be a real parameter to be chosen later and write $\mu^2(d) = M_Z(d) + R_Z(d)$ where

$$M_Z(d) = \sum_{\substack{l^2 | d \\ l \leq Z}} \mu(l) \quad \text{and} \quad R_Z(d) = \sum_{\substack{l^2 | d \\ l > Z}} \mu(l).$$

Define

$$S_M(X, Y; \hat{\phi}, f, \Phi) = \sum_{\gcd(d, 2)=1} M_Z(d) \sum_{p \leq Y} \frac{a_f(p) \log p}{\sqrt{p}} \left(\frac{8d}{p} \right) \hat{\phi} \left(\frac{\log p}{2 \log X} \right) \Phi \left(\frac{d}{X} \right)$$

and

$$S_R(X, Y; \hat{\phi}, f, \Phi) = \sum_{\gcd(d, 2)=1} R_Z(d) \sum_{p \leq Y} \frac{a_f(p) \log p}{\sqrt{p}} \left(\frac{8d}{p} \right) \hat{\phi} \left(\frac{\log p}{2 \log X} \right) \Phi \left(\frac{d}{X} \right),$$

so that $S(X, Y; \hat{\phi}, f, \Phi) = S_M(X, Y; \hat{\phi}, f, \Phi) + S_R(X, Y; \hat{\phi}, f, \Phi)$.

Using standard techniques (see (3.3)), we can show that by choosing Z appropriately that $S_R(X, Y; \hat{\phi}, f, \Phi)$ is small. Hence the main term arises only from $S_M(X, Y; \hat{\phi}, f, \Phi)$. We write it as

$$(2.7) \quad S_M(X, Y; \hat{\phi}, f, \Phi) = \sum_{p \leq Y} \frac{a_f(p) \log p}{\sqrt{p}} \left(\frac{8}{p} \right) \hat{\phi} \left(\frac{\log p}{2 \log X} \right) \left(\sum_{\gcd(d, 2)=1} M_Z(d) \left(\frac{d}{p} \right) \Phi \left(\frac{d}{X} \right) \right).$$

We now evaluate the inner sum above following a method of K. Soundararajan in [32] by applying the Poisson summation formula to the sum over d . For all odd integers k and all integers m , we introduce the Gauss-type sums

$$\tau_m(k) := \sum_{a \pmod{k}} \left(\frac{a}{k} \right) e \left(\frac{am}{k} \right) =: \left(\frac{1+i}{2} + \left(\frac{-1}{k} \right) \frac{1-i}{2} \right) G_m(k).$$

We quote Lemma 2.3 of [32] which determines $G_m(k)$.

Lemma 2.6. *If $(k_1, k_2) = 1$ then $G_m(k_1 k_2) = G_m(k_1) G_m(k_2)$. Suppose that p^a is the largest power of p dividing m (put $a = \infty$ if $m = 0$). Then for $b \geq 1$ we have*

$$G_m(p^b) = \begin{cases} 0 & \text{if } b \leq a \text{ is odd,} \\ \phi(p^b) & \text{if } b \leq a \text{ is even,} \\ -p^a & \text{if } b = a + 1 \text{ is even,} \\ (\frac{m/p^a}{p}) p^a \sqrt{p} & \text{if } b = a + 1 \text{ is odd,} \\ 0 & \text{if } b \geq a + 2. \end{cases}$$

For a Schwartz function F , we define

$$(2.8) \quad \tilde{F}(\xi) = \frac{1+i}{2} \hat{F}(\xi) + \frac{1-i}{2} \hat{F}(-\xi) = \int_{-\infty}^{\infty} (\cos(2\pi\xi x) + \sin(2\pi\xi x)) F(x) dx.$$

We quote Lemma 2.6 of [32] which determines the inner sum in (2.7).

Lemma 2.7. *Let Φ be a non-negative, smooth function supported in $(1, 2)$. For any odd integer k ,*

$$\sum_{\gcd(d, 2)=1} M_Z(d) \left(\frac{d}{k}\right) \Phi\left(\frac{d}{X}\right) = \frac{X}{2k} \left(\frac{2}{k}\right) \sum_{\substack{\alpha \leq Z \\ \gcd(\alpha, 2k)=1}} \frac{\mu(\alpha)}{\alpha^2} \sum_m (-1)^m G_m(k) \tilde{\Phi}\left(\frac{mX}{2\alpha^2 k}\right),$$

where $\tilde{\Phi}$ is as defined in (2.8).

Note that for any non-negative integer l ,

$$\tilde{\Phi}^{(l)}(\xi) \ll 1, \quad |\xi| < 1.$$

Also note via integration by parts,

$$\tilde{\Phi}(\xi) = \frac{-1}{2\pi\xi} \left(\int_1^{1+1/U} + \int_{2-1/U}^2 \right) \Phi'(x) (\sin(2\pi\xi x) - \cos(2\pi\xi x)) dx \ll \frac{1}{|\xi|}.$$

Similarly, one can show for any $l \geq 0, j \geq 1$,

$$\tilde{\Phi}^{(l)}(\xi) \ll \frac{U^{j-1}}{|\xi|^j}.$$

2.8. Primitive cubic and quartic Dirichlet characters. The classification of all the primitive cubic characters of conductor q coprime to 3 is given in [1]. It is shown there that every such character is of the form $m \rightarrow (\frac{m}{n})_3$ for some $n \in \mathbb{Z}[\omega]$, with $n \equiv 1 \pmod{3}$, n square-free and not divisible by any rational primes, $N(n) = q$. Here the symbol $(\frac{\cdot}{n})_3$ is the cubic residue symbol in the ring $\mathbb{Z}[\omega]$. For a prime $\pi \in \mathbb{Z}[\omega]$ with $N(\pi) \neq 3$, the cubic character is defined for $a \in \mathbb{Z}[\omega]$, $\gcd(a, \pi) = 1$ by $(\frac{a}{\pi})_3 \equiv a^{\frac{N(\pi)-1}{3}} \pmod{\pi}$, with $(\frac{a}{\pi})_3 \in \{1, \omega, \omega^2\}$. When $\pi|a$, set $(\frac{a}{\pi})_3 = 0$. One then extends the cubic character to composite n with $\gcd(N(n), 3) = 1$ multiplicatively.

Similarly, one can give a classification of all the primitive complex quartic characters of conductor q coprime to 2. Every such character is of the form $m \rightarrow (\frac{m}{n})_4$ for some $n \in \mathbb{Z}[i]$, with $n \equiv 1 \pmod{(1+i)^3}$, n square-free and not divisible by any rational primes and $N(n) = q$. Here the symbol $(\frac{\cdot}{n})_4$ is the quartic residue symbol in the ring $\mathbb{Z}[i]$. For a prime $\pi \in \mathbb{Z}[i]$ with $N(\pi) \neq 2$, the quartic character is defined for $a \in \mathbb{Z}[i]$, $\gcd(a, \pi) = 1$ by $(\frac{a}{\pi})_4 \equiv a^{\frac{N(\pi)-1}{4}} \pmod{\pi}$, with $(\frac{a}{\pi})_4 \in \{\pm 1, \pm i\}$. When $\pi|a$, $(\frac{a}{\pi})_4$ is defined to be zero. Then the quartic character can be extended to composite n with $\gcd(N(n), 2) = 1$ multiplicatively. Note that in $\mathbb{Z}[i]$, every ideal coprime to 2 has a unique generator congruent to 1 modulo $(1+i)^3$.

3. PROOF OF THEOREM 1.1

Note that as $X \rightarrow \infty$,

$$\sum_{d \in D(X)} \Phi\left(\frac{d}{X}\right) \sim \#D(X).$$

Moreover, as $\hat{\phi}$ is supported in $(-1, 1)$, we have

$$\int_{-\infty}^{\infty} \phi(t) dt + \frac{1}{2} \int_{-\infty}^{\infty} \hat{\phi}(u) du = \int_{-\infty}^{\infty} \phi(t) \left(1 + \frac{\sin(2\pi t)}{2\pi t}\right) dt.$$

Theorem 1.1 thus follows from (2.2), provided that we show for any Schwartz function ϕ with $\hat{\phi}$ supported in $(-1, 1)$,

$$(3.1) \quad \lim_{X \rightarrow \infty} \frac{S(X, Y; \hat{\phi}, f, \Phi)}{X \log X} = 0.$$

As

$$S(X, Y; \hat{\phi}, f, \Phi) = S_M(X, Y; \hat{\phi}, f, \Phi) + S_R(X, Y; \hat{\phi}, f, \Phi),$$

the remainder of this section is therefore devoted to the evaluation of $S_R(X, Y; \hat{\phi}, f, \Phi)$ and $S_M(X, Y; \hat{\phi}, f, \Phi)$.

3.1. Estimation of $S_R(X, Y; \hat{\phi}, f, \Phi)$. In this section, we estimate $S_R(X, Y; \hat{\phi}, f, \Phi)$. We first seek a bound for

$$E(Y; \chi, \hat{\phi}, f) := \sum_{p \leq Y} \frac{a_f(p) \log p}{\sqrt{p}} \chi(p) \hat{\phi}\left(\frac{\log p}{2 \log X}\right),$$

for any non-principal quadratic character χ with modulus q and $Y \leq X^{2-2\epsilon}$. For this we need the following result which follows from [18, Theorem 5.15].

Lemma 3.2. *Suppose that GRH is true. For any Dirichlet character χ with modulus q , we have for $x \geq 1$,*

$$\sum_{p \leq x} a_f(p) \chi(p) \log p \ll x^{1/2} \log^2(qx).$$

It follows from the above lemma and partial summation that

$$(3.2) \quad E(Y; \chi, \hat{\phi}, f) \ll \log^3(qX).$$

Now on writing $d = l^2 m$, we obtain

$$(3.3) \quad \begin{aligned} S_R(X, Y; \hat{\phi}, f, \Phi) &= \sum_{\substack{l > Z \\ \gcd(l, 2) = 1}} \mu(l) \sum_{\gcd(m, 2) = 1} \Phi\left(\frac{l^2 m}{X}\right) E(Y; \chi_{8l^2 m}, \hat{\phi}, f) \\ &\ll \sum_{l > Z} \sum_{X/l^2 \leq m \leq 2X/l^2} \log^3(X) \ll \frac{X \log^3 X}{Z}. \end{aligned}$$

3.3. Estimation of $S_M(X, Y; \hat{\phi}, f, \Phi)$. Applying Lemma 2.7 to the inner sum of (2.7), we see that the sum in $S_M(X, Y; \hat{\phi}, f, \Phi)$ corresponding to the contribution of $m = 0$ is zero, as it follows directly from the definition that $G_0(k) = \varphi(k)$ if k is a square and $G_0(k) = 0$ otherwise.

Now, the sums in $S_M(X, Y; \hat{\phi}, f, \Phi)$ corresponding to the contribution of $m \neq 0$ can be written as $XR/2$, where

$$R = \sum_{\substack{p \leq Y \\ \gcd(2, p) = 1}} \frac{a_f(p) \log p}{p} \hat{\phi}\left(\frac{\log p}{2 \log X}\right) \sum_{\substack{\alpha \leq Z \\ \gcd(\alpha, 2p) = 1}} \frac{\mu(\alpha)}{\alpha^2} \sum_{\substack{m = -\infty \\ m \neq 0}}^{\infty} \left(\frac{m}{p}\right) (-1)^m \tilde{\Phi}\left(\frac{mX}{2\alpha^2 p}\right).$$

We recast the condition $\gcd(p, 2\alpha) = 1$ as $\chi_{4\alpha^2}(p)$ and use estimation (3.2) to deduce by partial summation (note that we will take Z to be smaller than some power of X so that $\log \alpha \ll \log X$) that

$$\begin{aligned} \sum_{\substack{p \leq Y \\ \gcd(2\alpha, p)=1}} \frac{a_f(p) \log p}{p} \hat{\phi} \left(\frac{\log p}{2 \log X} \right) \tilde{\Phi} \left(\frac{mX}{2\alpha^2 p} \right) \left(\frac{m}{p} \right) &= \int_1^Y \frac{1}{\sqrt{V}} \tilde{\Phi} \left(\frac{mX}{2\alpha^2 V} \right) dE(V; \chi_{4\alpha^2 m}, \hat{\phi}, f) \\ &\ll \log^3(X(|m| + 2)) \left(\frac{1}{\sqrt{Y}} \left| \tilde{\Phi} \left(\frac{mX}{2\alpha^2 Y} \right) \right| + \int_1^Y \frac{1}{V^{3/2}} \left| \tilde{\Phi} \left(\frac{mX}{2\alpha^2 V} \right) \right| dV \right. \\ &\quad \left. + \int_1^Y \frac{X}{\alpha^2 V^{5/2}} \left| m \tilde{\Phi}' \left(\frac{mX}{2\alpha^2 V} \right) \right| dV \right). \end{aligned}$$

Hence we have

$$R \ll \sum_{\alpha \leq Z} \frac{1}{\alpha^2} (R_1 + R_2 + R_3),$$

where

$$R_1 = \frac{1}{\sqrt{Y}} \sum_{m \neq 0} \log^3(X(|m| + 2)) \left| \tilde{\Phi} \left(\frac{mX}{2\alpha^2 Y} \right) \right|, \quad R_2 = \int_1^Y \frac{1}{V^{3/2}} \sum_{m \neq 0} \log^3(X(|m| + 2)) \left| \tilde{\Phi} \left(\frac{mX}{2\alpha^2 V} \right) \right| dV$$

and

$$R_3 = \int_1^Y \frac{X}{\alpha^2 V^{5/2}} \sum_{m \neq 0} \log^3(X(|m| + 2)) \left| m \tilde{\Phi}' \left(\frac{mX}{2\alpha^2 V} \right) \right| dV.$$

We now gather the estimates in Appendix C of [26] for R_1, R_2 and R_3 (but be aware that the sum over p in [26] does not exist in our situation here):

$$(3.4) \quad R_1 + R_2 \ll \frac{U \alpha^2 \sqrt{Y} \log^7 X}{X} \text{ and } R_3 \ll \frac{U \alpha^2 \sqrt{Y} \log^7 X}{X} + \frac{U^2 \alpha^4 Y^{3/2} \log^7 X}{X^{2008}}.$$

Some of the estimates quoted above from [26] have their origins in [10]. The estimates from [26] would suffice for our purpose, but the improved estimates from [10] for $R_1 + R_2$ are neater to use here. Combining these estimates in (3.4), we obtain

$$R \ll \frac{U Z \sqrt{Y} \log^7 X}{X}.$$

Thus we conclude that the contribution of $m \neq 0$ is bounded by

$$(3.5) \quad U Z \sqrt{Y} \log^7 X.$$

3.4. Conclusion. We now combine the bounds (3.3), (3.5) and take $Y = X^{2-2\epsilon}, Z = \log^3 X$ (recall that $U = \log X$) with any fixed $\epsilon > 0$ to obtain

$$S(X, Y; \hat{\phi}, f, \Phi) \ll \frac{X \log^3 X}{Z} + U Z \sqrt{Y} \log^7 X = o(X \log X),$$

which implies (3.1) and this completes the proof of Theorem 1.1.

4. PROOF OF THEOREM 1.2

Let $N_d(X)$ be the number of primitive cubic characters of conductor $q \leq X$ with $\gcd(q, d) = 1$. It is shown in [5] that $N_d(X) \sim c(d)X$ as $X \rightarrow \infty$ for some constant $c(d)$. It follows from this that $\#C(X) \sim cX$ for some constant c as $X \rightarrow \infty$. Combining this with (2.1), we see that in order to establish Theorem 1.2, it suffices to show that for any Schwartz function f with $\hat{f}$ supported in $(-3/7, 3/7)$,

$$(4.1) \quad \lim_{X \rightarrow \infty} \frac{1}{X \log X} \sum_p \frac{\log p}{\sqrt{p}} \hat{f} \left(\frac{\log p}{\log X} \right) \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* (\chi(p) + \overline{\chi}(p)) = 0$$

and

$$(4.2) \quad \lim_{X \rightarrow \infty} \frac{1}{X \log X} \sum_p \frac{2 \log p}{p} \hat{f}\left(\frac{2 \log p}{\log X}\right) \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* (\chi(p^2) + \bar{\chi}(p^2)) = 0.$$

As both χ and $\bar{\chi}$ are primitive cubic characters, it is enough to consider the two limits for χ only. The term $p = 3$ in each sum above is $O(X)$. Hence we may assume $p \neq 3$ in the sums above and we apply the Cauchy-Schwarz inequality to see that

$$(4.3) \quad \begin{aligned} & \sum_{p \neq 3} \frac{\log p}{\sqrt{p}} \hat{f}\left(\frac{\log p}{\log X}\right) \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* \chi(p) \\ & \ll \left(\sum_{p \leq X^{1/5}} \frac{\log^2 p}{p} \right)^{1/2} \left(\sum_{3 \neq p \leq X^{1/5}} \left| \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* \chi(p) \right|^2 \right)^{1/2}, \end{aligned}$$

and

$$(4.4) \quad \begin{aligned} & \sum_{p \neq 3} \frac{\log p}{p} \hat{f}\left(\frac{2 \log p}{\log X}\right) \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* \chi(p^2) \\ & \ll \left(\sum_{p \leq X^{1/5}} \frac{\log^2 p}{p^2} \right)^{1/2} \left(\sum_{3 \neq p \leq X^{1/5}} \left| \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* \chi(p^2) \right|^2 \right)^{1/2}. \end{aligned}$$

It's easy to see that

$$\sum_{p \leq X^{1/5}} \frac{\log^2 p}{p} \ll \log^2 X \text{ and } \sum_{p \leq X^{1/5}} \frac{\log^2 p}{p^2} \ll 1.$$

Moreover, note that for a primitive cubic character χ , $\chi(p^2) = \bar{\chi}(p)$ which implies that the values of the sums on the right-hand sides of (4.3) and (4.4) are the same. Hence, it remains to estimate

$$(4.5) \quad \sum_{3 \neq p \leq Y} \left| \sum_{X \leq q \leq 2X} \sum_{\substack{\chi \pmod{q} \\ \chi^3 = \chi_0}}^* \chi(p) \right|^2.$$

Here, Y is a parameter independent of X . From our discussion in Section 2.8, we can recast (4.5) in terms of the cubic residue symbol as

$$\sum_{3 \neq p \leq Y} \left| \sum'_{\substack{N(n) \in \mathcal{I}(X) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n} \right)_3 \right|^2,$$

where the inner sum runs over square-free elements n of $\mathbb{Z}[\omega]$ that have no rational prime divisor and $\mathcal{I}(Z)$ henceforth denotes the dyadic interval

$$\mathcal{I}(Z) := [Z, 2Z]$$

for any real number Z . We now regard $(\frac{p}{\cdot})_3$ as a ray class group character ξ on $h_{(3p)}$ where we define $\xi((n)) = (\frac{p}{n})_3$. Now we remove the condition that n has no rational prime divisor by using Möbius inversion (note that one can uniquely express any $n \in \mathbb{Z}[\omega]$ as $n = n_1 n_2$, where $n_1 \in \mathbb{N}$, and n_2 has no rational prime

divisor), obtaining

$$\sum'_{\substack{N(n) \in \mathcal{I}(X) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 = \sum_{\substack{d \in \mathbb{Z}, d^2 \leq 2X \\ d \equiv 1 \pmod{3} \\ \gcd(d, p) = 1}} \mu_{\mathbb{N}}(|d|) \sum''_{\substack{N(n) \in \mathcal{I}(X/d^2) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3,$$

where the double prime indicates that nd is square-free (viewed as an element of $\mathbb{Z}[\omega]$). Here $\mu_{\mathbb{N}}$ is the usual Möbius function defined on $\mathbb{N}$. Note as d and p are coprime rational integers, it follows from the corollary to Proposition 9.3.4. of [16] that $(\frac{p}{d})_3 = 1$.

Since d is automatically square-free (as an element of $\mathbb{Z}[\omega]$), nd being square-free simply means that n is square-free and $\gcd(n, d) = 1$. Now use Möbius again (writing μ_{ω} for the Möbius function on $\mathbb{Z}[\omega]$) to detect the condition that n is square-free, getting

$$(4.6) \quad \sum'_{\substack{X \leq N(n) \leq 2X \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 = \sum_{\substack{d \in \mathbb{Z}, d^2 \leq 2X \\ d \equiv 1 \pmod{3} \\ \gcd(d, p) = 1}} \mu_{\mathbb{N}}(|d|) \sum_{\substack{N(l)^2 \leq 2X/d^2 \\ \gcd(l, d) = 1 \\ l \equiv 1 \pmod{3}}} \mu_{\omega}(l) \left(\frac{p}{l^2}\right)_3 \sum_{\substack{N(n) \in \mathcal{I}(X/(N(l)d)^2) \\ \gcd(n, d) = 1 \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3.$$

Here we changed variables via $n = l^2 n'$ and fixed l up to a unit by the condition $l \equiv 1 \pmod{3}$ (note that in $\mathbb{Z}[\omega]$, every ideal coprime to 3 has a unique generator congruent to 1 modulo 3).

We now apply a further Möbius to remove the condition $\gcd(n, d) = 1$ to get

$$(4.7) \quad \sum_{\substack{N(n) \in \mathcal{I}(X/(N(l)d)^2) \\ \gcd(n, d) = 1 \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 = \sum_{\substack{e|d \\ e \equiv 1 \pmod{3}}} \mu_{\omega}(e) \left(\frac{p}{e}\right)_3 \sum_{\substack{N(n) \in \mathcal{I}(X/N(e)N^2(l)d^2) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3.$$

Now we need the following lemma, which establishes a Pólya-Vinogradov type inequality for the cubic symbols:

Lemma 4.1. *Let $p \neq 3$ be a rational prime. Then we have*

$$(4.8) \quad \sum_{\substack{N(n) \leq X \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 \ll X^{1/3} p^{2/3} \log^2 p,$$

where the sum runs over elements $n \in \mathbb{Z}[\omega]$.

Proof. As we mentioned above, we regard $(\frac{p}{\cdot})_3$ as a ray class group character ξ on $h_{(3p)}$ so that we can recast the sum in (4.8) as

$$(4.9) \quad \sum_{\substack{N(n) \leq X \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 = \sum_{\substack{N(I) \leq X \\ \gcd(I, 3) = 1}} \xi(I),$$

where the sum above runs over non-zero integral ideals $I \in \mathbb{Z}[\omega]$. It's easy to see that ξ is induced by a primitive character of conductor (ap) for some $a|3$. Therefore, the condition $\gcd(I, 3) = 1$ imposed on the sum on the right-hand side of (4.9) implies that the said sum remains unchanged if ξ is replaced by ξ^* (say), the primitive character that induces ξ . We may therefore assume without loss of generality that ξ is primitive and we further use Möbius to detect the condition $\gcd(I, 3) = 1$ in the second sum in (4.9) while noting that the only ideals dividing 3 are (1) , $(1 - \omega)$ and (3) to get

$$\sum_{\substack{N(n) \leq X \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 = \sum_{\substack{N(I) \leq X \\ \gcd(I, 3) = 1}} \xi(I) = \sum_{h=1, 1-\omega, 3} \mu_{\omega}(h) \xi(h) \sum_{N(I) \leq X/N(h)} \xi(I).$$

Now we quote a result of E. Landau [23] (see also [33, Theorem 2]), which states that for an algebraic number field K of degree $n \geq 2$, ξ any primitive ideal character of K with conductor $\mathfrak{f}$, $k = |N(\mathfrak{f}) \cdot d_K|$ with d_K being the discriminant of K , we have for $X \geq 1$,

$$\sum_{N(I) \leq X} \xi(I) \leq k^{1/(n+1)} \log^n(k) \cdot X^{(n-1)/(n+1)},$$

where I runs over integral ideas of K .

We now identify $K = \mathbb{Q}(\omega)$ with $n = 2$ and $k = 3ap^2$, $a = 1, 3$ or 9 to see that the sum on the right-hand side of (4.9) is

$$O\left(X^{1/3}p^{2/3}\log^2 p\right).$$

This now completes the proof of the lemma. $\square$

Applying Lemma 4.1, we can majorize the left-hand side expression in (4.7) as

$$(4.10) \quad \sum_{\substack{N(n) \in \mathcal{I}(X/(N(l)d)^2) \\ \gcd(n,d)=1 \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 \ll \frac{X^{1/3}p^{2/3}\log^2 p}{(N(l)d)^{2/3}} \sum_{e|d} \frac{1}{\sqrt{N(e)}} \ll \frac{X^{1/3}p^{2/3}\log^2 p}{(N(l))^{2/3}d^{2/3-2\epsilon}},$$

for any $\epsilon > 0$. The last bound follows since we have $\#\{e \in \mathbb{Z}[\omega] : e|d\} \ll N(d)^\epsilon$. From (4.10) and (4.6), we have

$$\sum'_{\substack{N(n) \in \mathcal{I}(X) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 \ll X^{1/3}p^{2/3}\log^2 p \sum_{\substack{d \in \mathbb{N} \\ d^2 \leq 2X}} \frac{1}{d^{2/3-2\epsilon}} \sum_{\substack{N(l)^2 \leq 2X/d^2 \\ l \equiv 1 \pmod{3}}} \frac{1}{(N(l))^{2/3}}.$$

Note that it follows from [33, Theorem 2] that

$$\sum_{\substack{N(l) \leq X \\ l \equiv 1 \pmod{3}}} 1 \ll X.$$

We then deduce by partial summation that

$$\sum_{\substack{N(l)^2 \leq 2X/d^2 \\ l \equiv 1 \pmod{3}}} \frac{1}{(N(l))^{2/3}} \ll \left(\frac{X}{d^2}\right)^{1/6}.$$

Therefore

$$\sum'_{\substack{N(n) \in \mathcal{I}(X) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 \ll X^{1/2}p^{2/3}\log^2 p \sum_{\substack{d \in \mathbb{N} \\ d^2 \leq 2X}} \frac{1}{d^{1-2\epsilon}} \ll X^{1/2+3\epsilon}p^{2/3}\log^2 p.$$

Hence, we obtain

$$\sum_{3 \neq p \leq Y} \left| \sum'_{\substack{N(n) \in \mathcal{I}(X) \\ n \equiv 1 \pmod{3}}} \left(\frac{p}{n}\right)_3 \right|^2 \ll X^{1+6\epsilon} \sum_{p \leq Y} p^{4/3} \log^4 p \ll X^{1+6\epsilon} Y^{7/3} \log^3 Y.$$

Applying the above bound in the estimations (4.3) and (4.4), we find that (4.1) and (4.2) hold so long as $Y^{7/6} \leq X^{1/2-7\epsilon/2}$ and as ϵ is arbitrary, the proof of Theorem 1.2 is completed.

5. PROOF OF THEOREM 1.3 AND THEOREM 1.4

The proofs of both Theorems 1.3 and 1.4 are similar to that of Theorem 1.2 so we shall skip most of the details. For the proof of Theorem 1.3, one can show, similar to the proof of [11, Lemma 4.2] that

$$\#C_{(9)}(X) \sim c'X \text{ as } X \rightarrow \infty$$

for some constant c' . Recall that $C_{(9)}$ denotes the set of cubic symbols $\chi_c = \left(\frac{\cdot}{c}\right)_3$ with c square-free, congruent to 1 modulo 9 and $X \leq N(c) \leq 2X$. We then proceed as in the proof of Theorem 1.2 to see that it suffices to show for any fixed $\epsilon > 0$, we have

$$(5.1) \quad \sum_{\substack{\mathfrak{p} \\ 3 \neq N(\mathfrak{p}) \leq Y}} \left| \sum_{\substack{N(c) \in \mathcal{I}(X) \\ c \equiv 1 \pmod{9}}}^* \left(\frac{\mathfrak{p}}{c}\right)_3 \right|^2 \ll X^{1+\epsilon} Y^{5/3} \log^3 Y,$$

where Y is a parameter independent of X . We now regard $(\frac{\mathbf{p}}{c})_3$ as a ray class group character ξ on $h_{(3)\mathbf{p}}$ where we set $\xi((c)) = (\frac{\mathbf{p}}{c})_3$. Using the ray class characters on $h_{(3)\mathbf{p}}$ to detect the condition $c \equiv 1 \pmod{9}$ in the inner sum on the left-hand side of (5.1), we get

$$\sum_{\substack{N(c) \in \mathcal{I}(X) \\ c \equiv 1 \pmod{9}}}^* \left(\frac{\mathbf{p}}{c}\right)_3 = \frac{1}{\#h_{(9)}} \sum_{\psi \pmod{9}} \sum_{\substack{N(c) \in \mathcal{I}(X) \\ c \equiv 1 \pmod{3}}}^* \psi((c)) \xi((c)).$$

The estimation in (5.1) follows, after using Möbius to detect the condition that c is square-free, from the following estimation:

$$\sum_{\substack{N(c) \leq X \\ c \equiv l \pmod{9}}} \psi((c)) \xi((c)) \ll X^{1/3} N(\mathbf{p})^{1/3} \log^2 N(\mathbf{p})$$

where $l \equiv 1 \pmod{3} \in \mathbb{Z}[\omega]$. One more application of the ray class characters on $h_{(9)}$ shows that the above estimation follows from the following:

$$(5.2) \quad \sum_{\substack{N(I) \leq X \\ \gcd(I, 3) = 1}} \psi(I) \psi'(I) \xi(I) \ll X^{1/3} N(\mathbf{p})^{1/3} \log^2 N(\mathbf{p}),$$

where the sum above runs over non-zero integral ideals $I \in \mathbb{Z}[\omega]$. The character $\psi\psi'\xi$ can be viewed as a ray class group character on $h_{(9)\mathbf{p}}$ and our definition of ξ implies that it is induced from a primitive character on $h_{(a)\mathbf{p}}$ with $a|9$. Consequently, as the condition $\gcd(I, 3) = 1$ is imposed on the summation in (5.2), the value of this sum remains unaltered if $\psi\psi'\xi$ is replaced by the primitive character that induces it. We may therefore without loss of generality assuming $\psi\psi'\xi$ is primitive and an application of the Möbius function as in the proof of Lemma 4.1 allows us to obtain the desired bound in (5.2).

For the proof of Theorem 1.4, one can show, following the approach in [5], that

$$\#Q(X) \sim dX \text{ as } X \rightarrow \infty$$

for some constant d as $X \rightarrow \infty$. The rest of the proof goes in a similar fashion as that of Theorem 1.2.

6. NOTES

We remark here that it is conceivable that results along the lines of Theorems 1.2, 1.3 and 1.4 can be proved using a simpler approach involving mean-value estimates for sums of characters of a fixed order. This method was used in [26]. The afore-mentioned mean-value estimate for quadratic character sums is due to M. Jutila [20, Lemma 5], but the analogous results for characters of orders higher than two which would be needed here do not seem to be available. Moreover, if good mean value estimates can be obtained for cubic and quartic characters, one would expect that the support of $\hat{f}$ in Theorems 1.2, 1.3 and 1.4 can be significantly widened.

It would also be interesting to consider the one level density of low-lying zeros of families of Dirichlet L -function for characters of orders larger than 4. But the relation between higher order residue symbols and n -th order primitive Dirichlet characters would be more difficult to describe.

Acknowledgments. During this work, P. G. was supported by some postdoctoral research fellowships at Nanyang Technological University (NTU) and L. Z. by an Academic Research Fund Tier 1 Grant at NTU. Both authors wish to thank Stephan Baier for some interesting discussions concerning some parts of this paper and Matt Young for making them aware of the estimates in [23] which led to some improvements. Finally, the authors would also like to thank the referee of his/her many comments and suggestions.

REFERENCES

- [1] S. Baier and M. P. Young, *Mean values with cubic characters*, J. Number Theory (to appear). arXiv:0804.2233.
- [2] S. Baier and L. Zhao, *On the low-lying zeros of Hasse-Weil L -functions for elliptic curves*, Adv. Math. **219** (2008), no. 3, 952–985.
- [3] A. Brumer, *The average rank of elliptic curves, I*, Invent. Math. **109** (1992), no. 3, 445–472.

- [4] H. Davenport, *Multiplicative Number Theory*, Third Edition, Graduate Texts in Mathematics, vol. 74, Springer-Verlag, New York, 2000.
- [5] C. David, J. Fearnley, and H. Kisilevsky, *On the vanishing of twisted L -functions of elliptic curves*, Experiment. Math. **13** (2004), no. 2, 185–198.
- [6] P. Deligne, *La conjecture de Weil. I.*, Inst. Hautes Etudes Sci. Publ. Math. **43** (1974), 273–307.
- [7] E. Dueñez and S. J. Miller, *The low lying zeros of a $GL(4)$ and a $GL(6)$ family of L -functions*, Compos. Math. **142** (2006), no. 6, 1403–1425.
- [8] ———, *The effect of convolving families of L -functions on the underlying group symmetries*, Proc. London Math. Soc. (to appear). arXiv:math.NT/0607688.
- [9] E. Fouvry and H. Iwaniec, *Low-lying zeros of dihedral L -functions*, Duke Math. J. **116** (2003), no. 2, 189–217.
- [10] P. Gao, *n -level density of the low-lying zeros of quadratic Dirichlet L -functions*. arXiv:0806.4830.
- [11] A. M. Güloğlu, *On low lying zeros of automorphic L -functions*, Ph.D. Thesis, 2005. The Ohio State University.
- [12] ———, *Low-lying zeros of symmetric power L -functions*, Int. Math. Res. Not. **2005** (2005), no. 9, 517–550.
- [13] D. R. Heath-Brown, *The average rank of elliptic curves*, Duke Math. J. **122** (2004), no. 3, 225–320.
- [14] C. Hughes and S. J. Miller, *Low-lying zeros of L -functions with orthogonal symmetry*, Duke Math. J. **136** (2007), no. 1, 115–172.
- [15] C. P. Hughes and Z. Rudnick, *Linear statistics of low-lying zeros of L -functions*, Q. J. Math. **54** (2003), no. 3, 309–333.
- [16] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Second Edition, Graduate Texts in Mathematics, vol. 84, Springer-Verlag, New York, 1990.
- [17] H. Iwaniec, *Topics in Classical Automorphic Forms*, Graduate Studies in Mathematics, vol. 17, American Mathematical Society, Providence, 1997.
- [18] H. Iwaniec and E. Kowalski, *Analytic Number Theory*, American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, 2004.
- [19] H. Iwaniec, W. Luo, and P. Sarnak, *Low lying zeros of families of L -functions*, Inst. Hautes Etudes Sci. Publ. Math. **91** (2000), 55–131.
- [20] M. Jutila, *On the mean value of $L(1/2, \chi)$ for real characters*, Analysis **1** (1981), no. 2, 149–161.
- [21] N. Katz and P. Sarnak, *Random matrices, frobenius eigenvalues, and monodromy*, American Mathematical Society Colloquium Publications, vol. 45, American Mathematical Society, Providence, 1999.
- [22] ———, *Zeros of zeta functions and symmetries*, Bull. Amer. Math. Soc. **36** (1999), no. 1, 1–26.
- [23] E. Landau, *Verallgemeinerung eines Polyaschen Satzes auf algebraische Zahlkörper*, Göttinger Nachr. (1918), 478–488.
- [24] W. Luo, *On Hecke L -series associated with cubic characters*, Compos. Math. **140** (2004), no. 5, 1191–1196.
- [25] S. J. Miller, *One- and two-level densities for rational families of elliptic curves: evidence for the underlying group symmetries*, Compos. Math. **140** (2004), no. 4, 952–992.
- [26] ———, *A symplectic test of the L -functions ratios conjecture*, Int. Math. Res. Not. IMRN (2008). Art. ID rnm146, 36 pp.
- [27] A. E. Özlük and C. Snyder, *On the distribution of the nontrivial zeros of quadratic L -functions close to the real axis*, Acta Arith. **91** (1999), no. 3, 209–228.
- [28] G. Ricotta and E. Royer, *Statistics for low-lying zeros of symmetric power L -functions in the level aspect*, Forum Math. (to appear). arXiv:math.NT/0703760.
- [29] E. Royer, *Petits zéros de fonctions L de formes modulaires*, Acta Arith. **99** (2001), no. 2, 147–172.
- [30] M. O. Rubinstein, *Low-lying zeros of L -functions and random matrix theory*, Duke Math. J. **209** (2001), no. 1, 147–181.
- [31] Z. Rudnick and P. Sarnak, *Zeros of principal L -functions and random matrix theory*, Duke Math. J. **81** (1996), no. 2, 269–322.
- [32] K. Soundararajan, *Nonvanishing of quadratic Dirichlet L -functions at $s = \frac{1}{2}$* , Ann. of Math. (2) **152** (2000), no. 2, 447–488.
- [33] J. E. Sunley, *On the class numbers of totally imaginary quadratic extensions of totally real fields*, Bull. Amer. Math. Soc. **78** (1972), 74–76.
- [34] M. P. Young, *Low-lying zeros of families of elliptic curves*, J. Amer. Math. Soc. **19** (2005), no. 1, 205–250.

Div. of Math. Sci., School of Phys. & Math. Sci.,
 Nanyang Technological Univ., Singapore 637371
 Email: penggao@ntu.edu.sg

Div. of Math. Sci., School of Phys. & Math. Sci.,
 Nanyang Technological Univ., Singapore 637371
 Email: lzhao@pmail.ntu.edu.sg